

ROLE OF BIOTA IN MINERAL TRANSFORMATIONS AND TRANSPORT

Dr Steve Rogers

CRC LEME, CSIRO Land and Water, PMB 2, Glen Osmond SA 5064

steve.rogers@csiro.au

Microorganisms are likely to have played a significant role in the formation of the Australian regolith and in the transport and transformation of minerals within regolith materials. Single celled bacteria were the only forms of life on the planet for approximately 50% of earth's history, first appearing about 4000×10^6 years before present (BP). More complex eukaryotic organisms such as fungi first appeared around 2000×10^6 years BP. Fossil records suggest that these early single celled organisms had similar cell structure to modern bacteria. In contemporary regolith environments about 5000 microbial species have been isolated and identified, with one gram of surface regolith material containing between 10^6 - 10^9 cells. However, estimates of the total number of microorganisms in the regolith range from 100,000 to 1×10^6 species, suggesting up to 95% of regolith microorganisms are unknown to science.

There is much evidence to suggest a role for microbial processes in shaping contemporary regolith environments. For instance, the weathering of rocks as a result of microbial activity is thought to be the dominant form of chemical rock weathering in both geological and contemporary environments. Production of organic acids by bacteria (formic, acetic, lactic, pyruvic, succinic, 2-ketogluconic) and fungi (nitric, sulfuric, citric, oxalic, gluconic), is a biological mechanism these organisms have evolved in order to solubilise and obtain trace elements such as P and S from minerals.

Geochemical transformations resulting from microbial activity in the regolith are also well established. One of the most well known is the biological oxidation of pyritic and sulfide minerals, carried out by a group of bacteria known as chemolithoautotrophes. These organisms are highly adapted to low pH environments, and utilise Fe or S as electron donors in their respiratory chain, catalysing Fe and S oxidation through the production of iron or sulfur oxidase enzymes, bacteria are therefore the key biochemical catalysts of geochemical transformations in these environments. Bacteria responsible for the oxidation of sulfide minerals including As, Cu, Co, Fe, Ni, Mo, Pb, Zn have been isolated from sulfidic environments and historically classified as members of the *Thiobacillus* genus notably (*Thiobacillus* sp., *Acidothiobacillus* sp. *Sulfobacillus thermosulfidooxidans* and *Acidanus brierleyi*). (Mukhopadhyaya *et al.*, 2000).

Whilst biological processes responsible for mineral weathering and geochemical transformations have been identified we still lack a sound mechanistic understanding of the principle biological processes underpinning biogeochemical dispersion, mobilization, transport, and mineralisation of minerals & trace elements in the Australian regolith. A more important question is what is the relative role of biological and non-biological (abiotic) processes driving mineral transport, transformation and accumulation in the regolith.

BIOLOGICAL ACTIVITY AND BIOGEOCHEMICAL TRANSFORMATIONS

Much of the current approach to the study of microbial transformations of minerals has focussed on the identification of organisms, species diversity and population dynamics, (Baker and Banfield, 2003) and their relationship to rates of mineral transformation. In fact environmental microbiology research concerning defining the role of biota in driving biogeochemical transformations in general has focussed on establishing the relationship between bacterial diversity and rates of biogeochemical processes (Rogers and Colloff, 1999). In general there is a paucity of evidence from available studies linking changes in species diversity or shifts in microbial populations to changes in biogeochemical functionality. Traditional microbiological techniques applied to the study of regolith microbial populations fail to identify more than 5% of the species present. They also fail to determine the 'functional attributes' of microbial populations responsible for geochemical transformations.

FUNCTIONAL MOLECULAR TOOLS

An alternative approach to determining the mechanistic relationship between microbial activity and biogeochemical transformations is to target the 'molecular genetic' attributes of microbial populations as opposed to their phylogenetic attributes (Rogers et al. 2002). We have developed a new generation of molecular biology techniques to study the role of biota in mineral biogeochemical transformations, based on the direct extraction of nucleic acids (DNA and RNA) from regolith samples that remove the need to isolate/culture and identify individual organisms. They allow direct comparison of biogeochemical reaction kinetics, gene expression and biotic diversity of organisms, all in the same sample, and represent a major advance over traditional microbiology methods.

Examples of the application of molecular tools to the study of Regolith biogeochemical mineral transformations

i. Identification of chemolithotropic microbial functional genes involved in Sulfide mineral oxidation

The suite of genes encoding for sulphide mineral oxidation in bacteria has recently been identified. The *soxB* gene encodes for a diheme cytochrome c enzyme and has been shown to be essential for chemolithotropic sulfide mineral oxidation. This functional gene has also been identified in representatives of all known groups of chemolithotropic sulfur oxidising bacteria, making it an ideal candidate for our functional molecular approach. Polymerase chain reaction (PCR) techniques have been designed that amplify a 1000 base pair region of the *soxB* gene sequence (Petri et al. 2001), allowing us to apply our existing functional molecular techniques to the study of sulfur oxidising chemolithotropic bacteria activity and presence in sulfidic sediments.

In brief, DNA and RNA were extracted from sulfidic sediments using established techniques. Oligonucleotide primers were designed to amplify through PCR *soxB* gene sequences in sample DNA extracts. The presence of functional genes indicates the presence of the biological potential for S mineral oxidation reactions. Figure 1 summarises the detection of the *soxB* gene in sulfidic sediments actively oxidising in the River Murray Floodplain.




Figure 1 *soxB* functional gene DNA detected in extracts from sulfidic sediments . 1000 bp PCR product visualised on 2% agarose gel. **Lane 1** 100bp DNA ladder; **Lane 2-4** *A.caldus* *soxB* positive control 1000bp product; **Lanes 5-29** presence of *soxB* gene in sulfidic sediments; **Lane 30** DNA quantification marker

ii. Microbial ecology/population diversity of Bacteria Associated with Gold solubilisation and precipitation

A current LEME PhD project has identified the role of bacteria in gold precipitation and solubilisation. In order to identify microorganisms present on gold flakes, numbers of species and if the organisms are living, the molecular technique 16S rRNA Density Gradient Gel Electrophoresis (DGGE) analysis has been used (Muyzer, 1999). In brief a region of the bacterial 16S gene can be amplified in DNA extracts. PCR products are separated on a DGGE electrophoresis gel, that separates double stranded DNA. Each different bacterial species has a unique 16S gene sequence, therefore each band on the gel

represents an individual bacterial species. Individual sequences (bands) from the DGGE gel can be further analysed by sequence analysis in order to identify the species present. This technique allows the analysis of all organisms present, as no culturing and isolation of cells is required. Figure 2 shows a representative DGGE acrylamide gel showing bacterial species 16S gene diversity in DNA extracted from a single gold flake collected from the Tomakin Park gold mine. The successful extraction and amplification of DNA from single gold flakes, confirms that the microbial biofilm thought to be responsible for solubilisation and precipitation reactions is 'alive'.




Figure 2. 16S rRNA Density Gradient Gel Electrophoresis profiles of bacterial populations on gold flakes collected at the Tomakin park gold mine. 16S V3 region amplified in soil DNA extracts with degenerate oligonucleotide PCR primers 27FGC and 534R. PCR product visualised on a 35-60% urea/formamide acrylamide gel (8%), 120V 20hrs.

Both these CRC LEME sponsored studies have demonstrated the application of new molecular techniques to the study of biological mineral transformations in the regolith. The core aim of LEME activities in this area is the identification of mechanisms, processes and physico-chemical conditions in contemporary environments where biogenic weathering, biogeochemical transformations and dispersion dominate. Allowing the development of predictive models to identify where in the regolith biological mineral transformation and transport processes have occurred in the geological past.

Acknowledgements

Frank Reith CRC LEME PhD student for the gold flake DGGE figure, Naomi Grey for technical assistance with the sulfidic sediment *soxB* work

REFERENCES

- Baker B.J. & Banfield, J.F. 2003. Microbial communities in acid mine drainage. *FEMS Microbiology Ecology* **44**, 139-152.
- Mukhopadhyaya P.N., Deb C., Lahiri C. & Roy P. 2000. A *soxA* gene encoding a diheme cytochrome c and a *sox* locus, essential for sulfur oxidation in a new sulfur lithotrophic bacterium. *Journal of Bacteriology* **182**, 4278-4287.
- Muyzer G. 1999. DGGE/TGGE a method for identifying genes from natural ecosystems. *Current Opinion in Microbiology* **2**, 317-322.
- Petri R., Podgorsek L. & Imhoff J.F. 2001. Phylogeny and distribution of the *soxB* gene among thiosulfate-oxidising bacteria. *FEMS Microbiology Letters* **197**, 171-178
- Rogers S.L. & Colloff M. 1999. How functionally resilient are Australian production systems? Future concepts for the study of functional biology and functional resilience of soil systems. *Proceedings 'Fixing the Foundations', National symposium on the role of soil science in sustainable land and water management. 11-12 November 1999.* South Australian Research and Development Institute, Adelaide. Australian Academy of Science.
- Rogers S.L. Colloff M. & Gomez D. 2002. Detection and expression of *nifH* gene sequences in nucleic acid extracts from Australian agricultural soils. *Abstracts. 13th Australian Nitrogen Fixation Conference – Fixed Nitrogen in Sustainable Agricultural Systems.* September 24-27 Adelaide, South Australia.