

## ALUNITE AS A MINERAL HOST IN FERRICRETE FROM THE ENTERPRISE PIT, MOUNT GIBSON GOLD DEPOSIT.

Robert Hough<sup>1</sup>, Cajetan Phang<sup>1</sup>, Marc Norman<sup>2</sup> & Ravi Anand<sup>1</sup>

<sup>1</sup>CRC LEME, CSIRO Exploration and Mining, PO Box 1130, Bentley, WA, 6102

<sup>2</sup>Research School of Earth Sciences, Australian National University, ACT, 0200

### INTRODUCTION

The physical features of the landscape, bedrock geology and regolith profiles of the Mount Gibson gold deposit (300 km NNE of Perth) and its environs have all been studied extensively (e.g., Anand *et al.* 1989, Yeats & Groves 1998). Transported overburden at the Enterprise pit at Mt. Gibson has been subjected to ferruginisation, calcification and silicification. As previously reported (Hough *et al.* 2003) these sediments, and their weathering products, provide a good setting to study mineral and regolith features that act as hosts for ore-related elements (e.g., Au, Cu, Pb, As, Zn and As), and may provide pathfinders for underlying mineralisation. Ultimately, such studies are expected to identify minerals that act as deep geochemical sensors and are therefore, exploration sampling media in areas of deep cover.

### ENTERPRISE PIT

#### Transported overburden.

In the Enterprise pit *in situ* regolith is overlain by Tertiary and Quaternary sediments. Slabby to pisolitic ferricrete is formed in the Tertiary sediments with anomalous Au (up to 587 ppb), whereas white clay between the slabby layers contains 112 ppb Au and 85 ppm As. The uppermost sandy to sandy-clay Quaternary unit is 3-4 m thick, and has been calcified and silicified to produce calcrete and hardpans with 128-436 ppb Au.

Building on our previous study (Hough *et al.* 2003) we have conducted XRD analyses of the ferricrete. This revealed that it contains alunite, a K-Al sulphate that is commonly precipitated from acidic groundwaters, kaolinite, and only minor goethite. Alunite occurs within the matrix of the sample, and is very fine grained with a red-brown colour. It appears as a late stage, pervasive precipitate that surrounds clay spheres and clasts.

We have also performed new laser ablation ICP-MS (LA-ICP-MS) and electron microprobe studies on the ferricrete. LA-ICP-MS has been shown to be a powerful technique in the analysis of regolith samples, especially for Au (Le Gleuher 2003a, b, Hough *et al.* 2003). Our previous LA-ICP-MS analyses on the kaolinite-rich matrix from this ferricrete sample revealed that it contained 1-1.5 ppm Au, 290 ppm Cu and 360 ppm As. At this stage we interpreted that kaolinites within the matrix were authigenic, and as such were trapping hydromorphically dispersed Au from underlying mineralisation. Minor iron oxides in the matrix were thought to be preferential sinks for the Cu and As. Our new XRD data and some scanning electron microscopy work suggest this matrix is dominated by alunite.

An electron-microprobe transect from the core of a clay sphere into the alunite rich matrix reveals the sharp change in trace element concentrations with substantial Cu (290 ppm), Pb (500 ppm), and As (360 ppm) in the alunite-rich areas. The new LA-ICP-MS analyses confirm the probe data and, building on our preliminary results (Hough *et al.* 2003), several analyses found 1-2 ppm Au within alunite rich areas (Figure 1). Hematite veins and large kaolinite rich clasts within the ferricrete are devoid of Au and have much lower levels of Cu, Pb and As.

Butt (2001) reported a similar alunite-Au association in the mid-upper saprolite at the Mount Percy gold deposit near Kalgoorlie; bulk S contents of 0.5-5% were reported in this horizon.

During LA-ICP-MS spot analyses we were able to monitor the count levels for the Au isotope in the mass spectrometer as the laser gently ablated the samples. This allows us to reveal if the Au is present as small grains, whereby it would give peaks, or is homogeneously distributed giving a constant plateau. Within the alunite matrix Au counts formed a constant plateau suggesting the Au is homogeneously distributed throughout the ablated material, and is therefore likely to be either extremely fine grained or chemically bound within the alunite mineral lattice.


In the near future we hope to utilise X-ray adsorption spectroscopy using synchrotron radiation to investigate this further, and to attempt to identify the oxidation state of the Cu, Pb and As.

## SUMMARY

Minerals precipitated from groundwaters in the upper sediments of regolith profiles could provide an interesting sampling medium to search for hydromorphic dispersion from underlying, and deeply buried mineralisation. Alunite appears to be one such mineral occurring in the regolith profile of the Enterprise pit, Mount Gibson, that contains appreciable Au, Cu, Pb and As.

## REFERENCES

- ANAND R.R., SMITH R.E., INNES J. & CHURCHWARD H.M. 1989. Exploration geochemistry about the Mt. Gibson gold deposits, Western Australia. *CSIRO Division of Exploration Geoscience, Perth, Restricted Report 20R*, 96 pp. Reissued as *CRC LEME Open File Report 35*, 1998.
- BUTT C.R.M. 2001. Dispersion of gold and associated elements in the lateritic regolith, Mystery Zone, Mt Percy, Kalgoorlie, Western Australia. *Geochemistry, Exploration, Environment, Analysis* **1**, 291-306.
- HOUGH R.M., PHANG C & ANAND R.R. 2003. Minerals hosts for gold: A study of transported overburden at the Enterprise pit, Mount Gibson gold deposit. In: Roach I.C. ed. *Advances in Regolith*. CRC LEME, pp. 201-204.
- LE GLEUHER M. 2003a. Trace element-mineral associations in the regolith, Scuddles massive Cu-Zn sulphide deposit, Western Australia. *CRC LEME Restricted Report 195R*, 82 p.
- LE GLEUHER M. 2003b. Mineral hosts for gold and trace elements in the regolith, Boddington and Mt Percy Gold deposits, Western Australia. *CRC LEME Restricted Report 196R*, 95 p.
- YEATS C.J. & GROVES D.I. 1998. The Archaean Mount Gibson gold deposits, Yilgarn Craton, Western Australia: Products of combined synvolcanic and syntectonic alteration and mineralisation. *Ore Geology Reviews* **13**, 1-5, 103-129.



**Figure 1:** Optical micrograph of the alunite-rich ferricrete sample from the Enterprise Pit, Mount Gibson.