

CRC LEME

Open File Report 189

Regolith Geochronology and Landscape Evolution

Brad Pillans, Editor

Issued September 2006

Printed March 2007

INTRODUCTION

INTRODUCTION: THE LEME GEOCHRONOLOGY PROJECT

Brad Pillans

CRC LEME, Research School of Earth Sciences, ANU, Canberra, ACT, 0200

The current LEME Geochronology Project [full title: Geochronology and models of landscape evolution], grew out of the Paleomagnetic Dating Project that began in LEME 1 in 1995. Eleven years on, paleomagnetic dating remains at the forefront of the project, but is now supplemented by other dating methods including K/Ar and Ar/Ar, cosmogenic isotopes, luminescence, (U-Th)/He, U/Pb and U-series (Fig. 1). Of these, the latter three methods have yet to reach maturity and further work is required to fully test their applications to regolith materials; the other methods may be considered to be “well established”.

The papers in this volume arose from a Regolith Dating Methods Workshop, held in the Research School of Earth Sciences at the Australian National University on Wednesday 20th November 2002. Work being carried out within CRC LEME Geochronology Project was discussed at this workshop and is described in this volume.

Below I outline some of the highlights of the Geochronology Project and its forerunner, the Paleomagnetic Dating Project. Overviews of Australian regolith geochronology are given in Pillans (1998, 2005). An offshoot from the Geochronology Project, the History of Aridity Project is also briefly discussed.

LACHLAN FOLD BELT/WESTERN NEW SOUTH WALES

First results were obtained from Northparkes mine, where Eric Tonui was undertaking his PhD fieldwork (Tonui 1998). Oxidized saprolite yielded a surprisingly old paleomagnetic age of ~320 Ma (Pillans *et al.* 1999), which when combined with apatite fission track data, indicated episodes of kilometre-scale burial and denudation (O’Sullivan *et al.* 2000). Interestingly, recently published K/Ar ages from the Jenolan Caves (Osborne *et al.* 2006) support this regional story.

Further paleomagnetic dating has been undertaken by Martin Smith (2001; 2006) at McKinnons, Elura, New Cobar and Peak Hill mines, yielding generally younger (Cenozoic and Mesozoic) ages. As part of his PhD study, in western NSW, Martin Smith also undertook a study of the suitability of U/Pb dating of anatase in silcrete (See Smith *et al.* this volume).

YILGARN

An early focus in the Geochronology Project was paleomagnetic dating of regolith in the Yilgarn. Sampling was undertaken in open pit mines, which provided excellent, deep exposures. Initially it was hoped to be able to date the widespread, so called “lateritic residuum”; however, this generally proved unsuitable for paleomagnetic dating. On the other hand, oxidized saprolite, up to 100 m below the surface, turned out to be an ideal medium, and revealed a long history of weathering dating back, in some instances, to pre-Cenozoic times. Some early results were published in Anand & Paine (2002, Table 16). These and later results clearly showed that deep oxidation had preferentially occurred during two major episodes – around 60 Ma, and during the last 12 Ma. There is also evidence of an earlier deep oxidation episode at ~300-315 Ma at Laverton (Lancefield mine) and Meekatharra (St Barbara’s mine), and another at ~180 Ma at Kalgoorlie (Mt Percy). The result from Lancefield is particularly interesting because the oxidation occurs in a tillite which is overlain by shale of early Permian age (Eyles & de Broekert 2001). The paleomagnetic weathering age indicates that the tillite must be latest Carboniferous or older.

Recent studies (e.g. Pigeon *et al.* 2004; Heim *et al.* 2006) have demonstrated the potential for dating iron nodules in Western Australia using the (U-Th)/He method. Mark Paine (formerly Curtin University) and Martin Smith (ANU) have pioneered this technique in LEME.

Suzanne Simons and Alexander Nemchin have investigated U-series dating of opal veins at Yalgoo (Simons & Nemchin, this volume).

LUCAS HEIGHTS

In July 2002 I was asked to assist in determining the timing of last movement of two faults on the site of the replacement research reactor (RRR) at Lucas Heights in Sydney.

The steeply dipping faults could be traced for more than 100m across the RRR excavation. The eastern strand was a normal fault with a dip separation of 1-1.3 m, while the western strand was a reverse fault with a dip separation of 0.25-0.3 m. The key issue was to assess the seismic hazard posed by the faults.

With no record of historical record of seismicity or surface fault rupture, the only obvious constraint on the history of fault movement was that they offset Triassic age Hawksbury Sandstone. In other words at least one faulting event had occurred in the last 200 million years –hardly a close constraint. However, in the south wall of the excavation a layer of iron oxide was draped across, but not cut by the faults. Samples of the iron oxide layer were taken for paleomagnetic measurements, and showed that the layer had reverse polarity which must have been acquired prior to the last reversal of the Earth's magnetic field (the Matuyama/Bruhnes reversal dated at ~0.78 Ma (Pillans & Bourman 1996)). Thus the last fault movement occurred at least 0.78 million years ago. Indeed, the paleomagnetic pole position indicated an age of > 5 Ma (Pillans 2003), which was considered sufficiently long ago for the risk of future fault movement to be negligible, and for approval to be given to proceed with construction. The RRR will be commissioned in early 2007.

SOUTH AUSTRALIA

Following on from paleomagnetic studies of Pleistocene sections near Adelaide (Pillans & Bourman 1996, 2001), further paleomagnetic sampling was undertaken of deeply oxidized rocks in the Mount Lofty Ranges and at the Moonta copper mine on Yorke Peninsula. Ages from both areas were late Cenozoic (Figure 2), as are K/Ar ages on alunite clays from Moonta (J. Dunlap pers. comm. 2005).

Further north, a ferricrete-capped mesa near Marla has become something of a reference locality for the Geochronology Project – see Figure 2. Paleomagnetic samples from the ferricrete yield a well-defined pole with an age of ~60 Ma, indistinguishable from the regionally extensive Morney Profile in SW Queensland (Idnurm & Senior 1978). Preliminary (U-Th)/He ages (Smith 2006) on the ferricrete are slightly younger (~50 Ma), but are uncorrected for He diffusion losses.

At Beltana, in the Flinders Ranges, there has been much speculation about the origin of the willemite (Zn_2SiO_4) ore – is it supergene or hypogene? In 2001 I collected samples of coronadite associated with the willemite (Figure 3), which yielded a K/Ar age of 435 ± 5 Ma (Groves *et al.* 2003). The age of the coronadite, together with fluid inclusion studies that indicate temperatures of formation in the range 50-150°C, clearly support a hypogene origin (Groves *et al.* 2003). The Beltana results, along with new results from Trinity Well, are further discussed by Jim Dunlap in this volume.

NORTHERN TERRITORY

Prior to LEME, little was known about the age of regolith in the Northern Territory, with one notable exception - a K/Ar dating study of supergene manganese deposits on Groote Eylandt by Dammer *et al.* (1996). However, in 2003, Juan-Pablo Bernal completed a PhD study on the Ranger uranium mine (Bernal 2003), demonstrating that U-series dating was feasible on goethitic pisoliths less than 0.5 million years old (see Bernal *et al.* this volume).

Recently, two LEME projects (NT Regolith, and Tanami projects) provided the opportunity to carry out reconnaissance paleomagnetic dating on samples from a wide range of sites (Pillans & Craig 2005). Current detailed work in the Tanami region indicates that the deep oxidation extends back to ~320 Ma, and that the area has been continuously subaerially exposed for at least 100 Ma, and perhaps longer.

HISTORY OF ARIDITY PROJECT

This project, which ran from 2002 until 2006, was designed to investigate the chronology and evolution of aridity in Australia from about 30 million years ago to the present. Luminescence dating and cosmogenic nuclides indicate that dune systems and stony deserts (gibbers) are significantly older

than previously thought. Papers by Derek Fabel and John Chappell, in this volume, describe the theory and application of cosmogenic nuclide dating in the context of landscape evolution.

REFERENCES

- ANAND, R. & PAINE, M. 2002. Regolith geology of the Yilgarn Craton, Western Australia: implications for exploration. *Australian Journal of Earth Sciences* **49**, 3-162.
- BERNAL, J.P. 2003. *In situ measurement of U-series disequilibria in iron oxy/hydroxides and its application to weathering geochronology*. Unpublished PhD thesis, ANU, Canberra.
- DAMMER, D., MCDUGALL, I. & CHIVAS, A.R., 1996. Isotopic dating of supergene manganese oxides from the Groote Eylandt deposit. *Economic Geology* **91**, 386-401.
- EYLES, N. & DE BROEKERT, P. 2001. Glacial tunnel valleys in the Eastern Goldfields of Western Australia cut below the Late Paleozoic Pilbara ice sheet. *Palaeogeography, Palaeoclimatology, Palaeoecology* **171**, 29-40.
- GROVES, I.M., CARMAN, C.E. & DUNLAP, W.J., 2003. Geology of the Beltana willemite deposit, Flinders Ranges, South Australia. *Economic Geology* **98**, 797-818.
- HEIM, J.A., VASCONCELOS, P.M., SHUSTER, D.L., FARLEY, K.A. & BROADBENT, G. 2006. Dating paleochannel iron ore by (U-Th)/He analysis of supergene goethite, Hamersley province, Australia. *Geology* **34**, 173-176.
- IDNURM, M. & SENIOR, B.R. 1978. Palaeomagnetic ages of Late Cretaceous and Tertiary weathered profiles in the Eromanga Basin, Queensland. *Palaeogeography, Palaeoclimatology, Palaeoecology* **24**, 263-277.
- OSBORNE, R.A.L., ZWINGMANN, H., POGSON, R.E. & COLCHESTER, D.M., 2006. Carboniferous clay deposits from Jenolan Caves, New South Wales: implications for timing of speleogenesis and regional geology. *Australian Journal of Earth Sciences* **53**, 377-405.
- O'SULLIVAN, P.B., GIBSON, D.L., KOHN, B.P., PILLANS, B. & PAIN, C.F. 2000. Long-term landscape evolution of the Northparkes region of the Lachlan Fold Belt, Australia: Constraints from fission track and paleomagnetic data. *Journal of Geology* **108**, 1-16.
- PIDGEON, R.T., BRANDER, T. & LIPPOLT, H.J, 2004. Late Miocene (U+Th)-⁴He ages of ferruginous nodules from lateritic duricrust, Darling Range, Western Australia. *Australian Journal of Earth Sciences* **51** 901-909.
- PILLANS, B. 1998. *Regolith Dating Methods. A Guide to Numerical Dating Techniques*. Cooperative Research Centre for Landscape Evolution and Mineral Exploration, Perth, 30 pp.
- PILLANS, B. 2002. Climate-driven weathering episodes during the last 200 Ma in Southern Australia. In: Preiss, V.P. ed. *Geoscience 2002: Expanding Horizons. Abstracts of the 16th Australian Geological Convention. Geological Society of Australia, Adelaide Convention Centre, Adelaide, SA, Australia. July 1-5 2002*, pp. 428.
- PILLANS, B. 2003. Dating ferruginous regolith to determine fault capability at Lucas Heights, Sydney. In: Roach, I.C. ed. *Advances in Regolith: Proceedings of the CRC LEME Regional Regolith Symposia, 2003*, 324-327.
- PILLANS, B. 2005. Geochronology of the Australian regolith. In: Anand, R.R. & de Broekert, P eds. *Regolith-Landscape Evolution Across Australia*. CRC LEME Monograph, pp. 41-61.
- PILLANS, B. & BOURMAN, R., 1996. The Brunhes/Matuyama polarity transition (0.78 Ma) as a chronostratigraphic marker in Australian regolith studies. *AGSO Journal of Australian Geology and Geophysics* **16(3)**, 289-294.
- PILLANS, B. & BOURMAN, R., 2001. Mid Pleistocene arid shift in southern Australia, dated by magnetostratigraphy. *Australian Journal of Soil Research* **39**, 89-98.

- PILLANS, B. & CRAIG, M. 2005. Reconnaissance paleomagnetic dating of regolith samples from the Northern Territory. *In: Roach, I.C. ed. Regolith 2005. Ten Years of CRC LEME.* CRC LEME, pp. 248-251.
- PILLANS, B., TONUI, E. & IDNURM, M. 1999. Paleomagnetic dating of weathered regolith at Northparkes mine, NSW. *In: Taylor, G. & Pain, C. eds. New Approaches to an Old Continent.* Cooperative Research Centre for Landscape Evolution and Mineral Exploration, Perth, pp. 237-242.
- SCHMIDT, P.W. & EMBLETON, B.J.J. 1976. Palaeomagnetic results from sediments of the Perth Basin, Western Australia, and their bearing on the timing of regional lateritisation. *Palaeogeography, Palaeoclimatology, Palaeoecology* **19**, 257-273.
- SMITH, M. 2001. *Palaeomagnetic age and geochemical characteristics of the regolith in the Cobar Basin, NSW.* Unpublished BSc (Hons) thesis, University of New South Wales, Sydney.
- SMITH, M. 2006. *Towards a geochronology of long term landscape evolution in northwestern NSW.* Unpublished PhD thesis, ANU, Canberra.
- TONUI, E. 1998. *Regolith mineralogy and geochemistry at Goonumbla, Parks, NSW.* Unpublished PhD thesis, ANU, Canberra.